

ChemBioDefense November, 2003

In vitro Characterization of Recombinant **Human BuChE** (Protexia) as a **Potential** Bioscavenger

David E. Lenz, Ph.D.,
Pharmacology Division
U.S. Army Medical Research
Institute of Chemical Defense
Aberdeen Proving Ground, MD
21010

USAMRICD

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to completing and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar DMB control number.	ion of information. Send comments arters Services, Directorate for Infor	regarding this burden estimate of mation Operations and Reports	or any other aspect of th s, 1215 Jefferson Davis I	is collection of information, Highway, Suite 1204, Arlington		
1. REPORT DATE 01 OCT 2005		2. REPORT TYPE N/A		3. DATES COVE	RED		
4. TITLE AND SUBTITLE					5a. CONTRACT NUMBER		
In vitro Characterization of Recombinant Human BuChE (Protexia) as a Potential Bioscavenger					5b. GRANT NUMBER		
Potential bioscavei	nger	5c. PROGRAM ELEMENT NUMBER					
6. AUTHOR(S)					5d. PROJECT NUMBER		
					5e. TASK NUMBER		
		5f. WORK UNIT NUMBER					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Pharmacology Division U.S. Army Medical Research Institute of Chemical Defense Aberdeen Proving Ground, MD 21010					8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)					10. SPONSOR/MONITOR'S ACRONYM(S)		
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)					
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release, distributi	on unlimited					
	otes 51, Proceedings of t Research, 17-20 No						
14. ABSTRACT							
15. SUBJECT TERMS							
16. SECURITY CLASSIFICATION OF: 17. LIN				18. NUMBER	19a. NAME OF		
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	ABSTRACT UU	OF PAGES 20	RESPONSIBLE PERSON		

Report Documentation Page

Form Approved OMB No. 0704-0188

Co-Authors

D.M. Cerasoli¹, E.M. Griffiths¹, B. P. Doctor², A. Saxena², N.H. Greig³, Q.S. Yu³, .Y. Huang⁴, H. Wilgus⁴ and C. N. Karatzas.⁴


¹Pharmacology Division, USAMRICD, 3100 Ricketts Pt. Rd, APG, MD 21010

- ² Biochemistry Division, 503 Robert Grant Dr., WRAIR, Bethesda, MD 20910
- ³ Drug Design & Development Section, NIA, 5600 Nathan Shock Drive, Baltimore, MD 21224
- ⁴ Nexia Biotechnologies Inc, 1000 St. Charles Ave, Block B, Vaudreuil-Dorion, Quebec, J7V8P5 Canada



Objectives

- Demonstrate improved medical protection against nerve agents
- Develop a prophylactic that detoxifies nerve agents at a rate sufficient to protect against 5LD₅₀ exposure
- Prophylactic should:
 - Be non-toxic
 - Produce no adverse side effects
 - Have no adverse effect on performance
 - Be easy to administer
 - Have a long biological half-life



3-D model of BuChE allows for rational drug design and molecular biology approaches to development of biological scavengers.



Concept

◆Provide a pretreatment capable of protecting against up to 5xLD₅₀ of nerve agent with no side effects and no need for additional protective clothing or therapy



Potential Bioscavengers


Efforts to date have focused on BuChE of human origin

- Production of Hu BChE From Cohn Fraction IV-4
 - •single band on SDS-PAGE
 - •~ 20 g =14 million units of purified BChE
 - •>98% pure; specific activity ~700 U/mg
- •Material (ProtexiaTM) from milk of transgenic goats is now available
 - •Hu-BuChE gene fused to a milk promoter
 - •Expression is directed to the milk of transgenic goats
 - •Purified material has Specific activity ~595 U/mg

Need to characterize ProtexiaTM versus plasma derived material




ProtexiaTM Binds H³-Soman






ProtexiaTM Binds H³-DFP





ProtexiaTM BuChE is Inhibited by OPs






Titration of h-BuChE and Protexia[™] by Soman and VX







Structures of Inhibitors






Structures of Inhibitors




Tolserine with h-BuChE






Tolserine with h-BuChE






Tolserine with RHu-BuChE (ProtexiaTM)






Tolserine with Protexia™






Phenserine with h-BuChE





Phenserine with ProtexiaTM





K_i and IC50 data for Set of Inhibitors

Inhibitor compound	Protexia™		Hu-BuChE	
	IC 50	<u>K_i (M)</u>	<u>IC 50</u>	<u>K_i (M)</u>
Phenserine	5 x 10 ⁻⁷	1.3 x 10 ⁻⁶	5 x 10 ⁻⁷	6.6 x 10 ⁻⁷
Tolserine	8 x 10 ⁻⁸	1.4 x 10 ⁻⁷	5 x 10 ⁻⁸	7.3 x 10 ⁻⁸
Phenethyl cynserine	5 x10 ⁻¹¹	5.1 x 10 ⁻¹⁰	1x 10 ⁻¹⁰	1.8 x 10 ⁻⁸
Ethyl carbamoyl	5 x 10 ⁻¹¹	6.2 x 10 ⁻¹⁰	5 x 10 ⁻¹⁴	1.9 x 10 ⁻¹³
Pyridostigmine bromide	5 x 10 ⁻⁷	4.6 x 10 ⁻⁷	3 x 10 ⁻⁷	6.2 x 10 ⁻⁷



Comparison of Hu-BuChE and ProtexiaTM

- In vitro comparison of both forms of human BuChE with a variety of inhibitors
- •Properties of recombinant HuBuChE from milk of transgenic goats (ProtexiaTM)
 - •Binds the nerve agents GA, GB, GD, VX
 - •In vitro properties similar to human plasma BuChE
- Results to date support similarity in the two sources.



Summary

Plasma BuChE is current standard.

- Provides protection
- •PK known in three species

ProtexiaTM binds all nerve agents

- •Binding is active site specific
- •Reaction with a variety of inhibitors is the same for both types of BuChE

While further characterization of Protexia $^{\rm TM}$ is needed to include PK and efficacy data, in vitro tests suggest both forms of the enzyme behave in a similar manner.

